

COSAVET-DF EDGE[®]



PROVEN
ECONOMICAL
CONVENIENT



COSAVET DF FUNGICIDE

COSAVET is a dry flowable sulphur fungicide that prevents powdery mildew and controls erinium mite of grape. Its patented formulation ensures a low dust, easy to mix product that helps to minimize the risk of scorching

Variations in particle size ensure immediate, mid-term, and residual activity through contact and vapour action to protect against target fungi.



TECHNICAL INFORMATION

For use on: Grapes, Apples, Pears, Stonefruit, Saskatoon berries, Greenhouse Cucumbers, Peas

Active Ingredients: Sulphur


Formulation: Dry Flowable

Packaging: 13.6 kg bag

Mode of Action: Disrupts cellular function by entering fungal cells and converting to hydrogen sulphide.

FRAC Group: M

Key diseases controlled: Powdery mildew, Brown rot, Leaf spot, Scab, Rust mite, Grape erineum mite, Entomosporium leaf spot, Entomosporium berry spot.



COSAVET DF rapidly disperses in water and stays in suspension, ensuring that all the COSAVET DF put in spray tank reaches your crop.

CROP	DISEASE CONTROLLED	RECOMMENDATIONS	RATE
See label for PHI			
APPLES PEARS	Powdery mildew Scab	Apply in regular protective schedule from green tip up to and including first cover spray.	750 g / 100L (22.5 kg/Ha)
		Apply in later cover sprays, and adjust rate depending on mildew severity and temperature.	400 – 500 g / 100 L water (12 – 15 kg. /Ha)
	Rust mites	Apply in summer sprays.	200 g / 100 L water (6.0 kg / Ha)
GRAPES	Powdery mildew	Apply at first sign of mildew and repeat after 10 days	420 g / 100L (12.6 kg. / Ha)
	Grape erineum mite	Apply at pre-bloom immediately after detection of erineum development on the leaves and again at mid-season.	3.4 kg. / Ha
PEACHES	Brown rot Leaf spot Scab	Apply in pink, bloom, shuck, cover and pre-harvest spray. Apply just before picking and repeat between picks if weather is wet.	750 g / 100L (22.5 kg. / Ha)
	Powdery mildew	Apply at husk-fall and repeat in 10 – 14 days	750 g / 100L (22.5 kg. / Ha)
PLUMS	Brown rot Leaf spot	Apply in pink, bloom, shuck, cover and pre-harvest spray. Apply just before picking and repeat between picks if weather is wet.	750 g / 100L (22.5 kg. / Ha)
SWEET CHERRIES	Brown rot	Apply in first, second and third cover sprays.	750 g / 100L (22.5 kg. / Ha)
SOUR CHERRIES	Brown rot	Apply at bloom stage only.	750 g / 100L (22.5 kg. / Ha)
	Powdery mildew	Apply in first, second and third cover sprays	400 g / 100L (12 kg. / Ha)
	Rust mites	Apply in summer sprays.	200 g / 100L (6 kg. / Ha)
SASKATOON BERRIES	Entomosporium leaf spot Entomosporium berry spot	Apply first at flower bud break and at 10 – 14 day intervals while disease risk persists.	7.5 kg. / Ha
CUCUMBERS (greenhouse)	Powdery mildew	Apply once every 5 days as required.	120 g / 100 L water
PEAS	Powdery mildew	Spray at first appearance of disease and repeat at 7 – 10 day intervals as necessary.	1.5 kg. / Ha
Do not use on Concord, Foch or DeChanac grape varieties (see label)			
Always read and follow label directions.		Cosavet is a registered trademark of Sulfur Mills Ltd.	