

2022-07-28
2021-2497

GROUP 50 FUNGICIDE

Container

PROPERTY® 300SC Fungicide

**For control or suppression of powdery mildew on outdoor
ornamentals and labeled crops.**

COMMERCIAL

SUSPENSION

READ THE LABEL AND ATTACHED PAMPHLET BEFORE USING

ACTIVE INGREDIENT: PYRIOFENONE..... 300 g/L

Contains 1,2-benzisothiazolin-3-one as a preservative at 0.035% as a preservative
OR

Contains 2-Bromo-2-nitropropane-1,3-diol at 0.0088%
and

5-Chloro-2-methyl-4-isothiazolin-3-one at 0.00084%
and

2-Methyl-4-isothiazolin-3-one at 0.00028% as preservatives

REGISTRATION NO: 32534 PEST CONTROL PRODUCTS ACT

Net Contents: 500 ml to 200 L

Registrant
ISK Biosciences Corporation
7470 Auburn Road, Suite A
Concord, Ohio
44077
USA
1-877-706-4640

Distributed By:
Belchim Crop Protection
104 Cooper Drive, Unit 3
Guelph, ON
N1C 0A4
Canada
866-613-3336

Property is a registered trademark of Ishihara Sangyo Kaisha, Ltd.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

DO NOT apply by air.

DO NOT apply in greenhouses.

Not for use in Residential Areas.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab.

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

FIRST AID

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

In Case of Emergency: Call 1-888-484-7546.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms. Observe buffer zones specified under **DIRECTIONS FOR USE**.

Pyriofenone is persistent and may carryover. It is recommended that any products containing pyriofenone not be used in areas treated with this product during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

STORAGE

To prevent contamination store this product away from food or feed.

DISPOSAL

For recyclable containers:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For returnable containers:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For containers that can be refilled for the user by the distributor/dealer:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Pamphlet

PROPERTY® 300SC Fungicide

**For control or suppression of powdery mildew on outdoor
ornamentals and labeled crops.**

COMMERCIAL

SUSPENSION

READ THE LABEL AND ATTACHED PAMPHLET BEFORE USING

ACTIVE INGREDIENT: PYRIOFENONE..... 300 g/L

Contains 1,2-benzisothiazolin-3-one as a preservative at 0.035% as a preservative

OR

Contains 2-Bromo-2-nitropropane-1,3-diol at 0.0088%

and

5-Chloro-2-methyl-4-isothiazolin-3-one at 0.00084%

and

2-Methyl-4-isothiazolin-3-one at 0.00028% as preservatives

REGISTRATION NO: 32534 PEST CONTROL PRODUCTS ACT

Net Contents: 500 ml to 200 L

Registrant
ISK Biosciences Corporation
7470 Auburn Road, Suite A
Concord, Ohio
44077
USA
1-877-706-4640

Distributed By:
Belchim Crop Protection
104 Cooper Drive, Unit 3
Guelph, ON
N1C 0A4
Canada
866-613-3336

Property is a registered trademark of Ishihara Sangyo Kaisha, Ltd.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

DO NOT apply by air.

DO NOT apply in greenhouses.

Not for use in Residential Areas.

For all crops, do not enter or allow worker entry into treated area during the restricted entry interval (REI) of 12 hours.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab.

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

FIRST AID

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

In Case of Emergency: Call 1-888-484-7546.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms. Observe buffer zones specified under DIRECTIONS FOR USE.

Pyriofenone is persistent and may carryover. It is recommended that any products containing pyriofenone not be used in areas treated with this product during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

STORAGE

To prevent contamination store this product away from food or feed.

DISPOSAL

For recyclable containers:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For returnable containers:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For containers that can be refilled for the user by the distributor/dealer:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

DIRECTIONS FOR USE

PROPERTY 300SC Fungicide is a fungicide with preventative and locally systemic properties for foliar diseases. PROPERTY 300SC Fungicide must be applied in scheduled protective programs and used in rotation with products with a different mode of action.

Under conditions favorable for disease development, the highest rate specified and shortest application interval should be used. Thorough, uniform coverage is essential for disease control.

MIXING

NOTE: Slowly invert container several times to assure uniform mixture of formulation before adding this product to the spray tank.

The required amount of PROPERTY 300SC Fungicide should be added slowly into the spray tank during filling. With concentrate sprays, pre-mix the required amount of PROPERTY 300SC Fungicide in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations. DO NOT allow spray mixture to stand overnight or for prolonged periods. Prepare only the amount of spray required for immediate use. Spraying equipment should be thoroughly cleaned immediately after the application.

TANK MIX COMPATIBILITY

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact ISK Biosciences Corporation at 1-877-706-4640 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label.

APPLICATION

Apply PROPERTY 300SC Fungicide in sufficient water to obtain adequate coverage of the foliage. Water volume to be used will vary with crop and amount of plant growth. Spray volume will usually range from 200 to 1000 litres per hectare for dilute ground sprays. Do not apply by air. Not for use in greenhouses.

DIRECTIONS FOR USE			
Crop	Diseases	Use Rate L Product Per Hectare	Instructions
Cucurbits Vegetables Crop Group 9	Powdery Mildew (<i>Podosphaera xanthii</i> , <i>Erysiphe cichoracearum</i>)	0.3 to 0.366 (90 to 110 g a.i./ha)	Application Instructions: Initiate applications prior to disease development and continue on a 7 to 10 day interval beginning with initial flowering or

			<p>when disease conditions are favorable for disease development, but prior to disease development. Use the low rate and long interval as disease preventative sprays or when disease conditions are low. Increase to highest rate and shortest interval under moderate to heavy disease pressure. Normal spray volumes range from 200 to 1000 litres per hectare for dilute ground sprays</p> <p>Resistance Management: Do not make more than 2 consecutive applications of PROPERTY 300SC Fungicide or other Group 50 containing fungicides before rotating to a fungicide with a different mode of action.</p> <p>Restrictions: Do not apply more than 1.2 L/ha/year The Pre-Harvest Interval (PHI) for this crop is 0 days.</p>
<p>Includes all members of the Cucurbit Vegetable Crop Group 9: Chayote (fruit); Chinese waxgourd (Chinese preserving melon); citron melon; cucumbers; gherkin; gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); <i>Momordica spp</i> (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); muskmelon, <i>Cucumis melo</i> (includes true cantaloupe, cantaloupe, casaba, Santa Claus melon; crenshaw melon; honeydew melon, honey balls, Persian melon, golden pershaw melon, mango melon, pineapple melon, snake melon); pumpkin; squash, summer (<i>Cucurbita pepo</i> includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash; watermelon (<i>Citrullus spp.</i>); and hybrids and/or varieties of these.</p>			
Grapes	Powdery mildew (<i>Erysiphe necator</i>)	0.3 to 0.366 (90 to 110 g a.i./ha)	<p>Application Instructions: For powdery mildew control, begin fungicide applications preventatively and continue as needed as per instructions below. Use the low rate and long interval as disease preventative sprays or when disease conditions are low. Increase to highest rate and shortest interval under moderate to heavy disease pressure. Normal spray volumes range from 200 to 1000 litres per hectare for dilute ground sprays, but can be up to 2000 litres per hectare for vineyards.</p> <p>Grapes: For use on all types of grapes (wine, table, raisin, and juice). Applications</p>
Strawberries	Powdery mildew (Suppression) (<i>Podosphaera aphanis</i>)		
Caneberries (blackberry; loganberry; red and black raspberry; wild raspberry;	Powdery mildew (Suppression) (<i>Sphaerotheca macularis</i>)		

cultivars and/or hybrids of these)			to grape should be made on a 14-day interval.
Goose Berries	Powdery mildew (Suppression) (<i>Sphaerotheca macularis</i>)		Applications to strawberries, caneberries, goose berries and Saskatoon berries should be made on a 7 to 10-day interval.
Saskatoon Berries	Powdery mildew (Suppression) (<i>Sphaerotheca macularis</i>)		<p>Resistance Management: Do not make more than 2 consecutive applications of PROPERTY 300SC Fungicide or other Group 50 containing fungicides before rotating to a fungicide with a different mode of action.</p> <p>Restrictions: Do not apply more than 1.2 L/ha/year. The Pre-Harvest Interval (PHI) for this crop is 0 days.</p>

DIRECTIONS FOR USE			
Crop	Diseases	Use Rate L Product Per Hectare	Instructions
Tomatoes	Powdery mildew (Suppression) (<i>Leveillula taurica</i>)	0.3 to 0.366 (90 to 110 g a.i./ha)	<p>Apply prior to disease initiation on a 7 to 14-day schedule when conditions are favorable for disease development.</p> <p>Use the low rate and long interval when conditions favor disease development and disease pressure is low. Increase use to the highest rate and shortest interval under heavy disease pressure. Normal spray volumes range from 200 to 1000 litres per hectare for dilute ground sprays.</p> <p>Resistance Management: Do not make more than 2 sequential applications of PROPERTY 300SC or other Group 50 containing fungicides before rotating to a fungicide with a different mode of action.</p> <p>Restrictions: Do not apply more than 1.2 L/ha/year The Pre-Harvest Interval (PHI) for this crop is 0 days.</p>

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than ISK Biosciences Corporation under the User Requested Minor Use Label Expansion program. For these uses, ISK Biosciences Corporation has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

For Application to Ornamentals, Grown Outdoors

DIRECTIONS FOR USE			
Crops	Diseases	Use Rate L Product Per Hectare	Instructions
Outdoor Ornamental Plants <i>Rosa</i> spp., <i>Photinia</i> spp., and non-bearing <i>Prunus</i> spp.	Powdery mildew (<i>Podosphaera</i> spp.) Powdery mildew (<i>Sphaerotheca</i> <i>pannosa</i>)	0.3 to 0.366 (90 to 110 g a.i./ha)	Application Instructions: Initiate applications prior to disease development and continue on a 7 to 10 day interval beginning when disease conditions are favorable for disease development, but prior to disease development. Use the low rate and long interval as disease preventative sprays or when conditions favour low disease pressure. Increase to highest rate and shortest interval under conditions conducive to moderate to heavy disease pressure. Normal spray volumes range from 200 to 1000 litres per hectare for dilute ground sprays Resistance Management: Do not make more than 2 consecutive applications of PROPERTY 300SC Fungicide or other Group 50 containing fungicides before rotating to a fungicide with a different mode of action. Restrictions: Do not apply more than 1.2 L/ha/year. Maximum of 3 applications at the high rate and a maximum of 4 applications at the low rate.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Airblast application: DO NOT apply during periods of dead calm. Avoid application of this

product when winds are gusty. DO NOT direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. DO NOT apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

DO NOT apply by air.

Buffer zones:

Use of the following spray methods or equipment do not require a buffer zone: hand-held or backpack sprayer and spot treatment.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Buffer Zones (metres) Required for the Protection of:			
			Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:	
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m
Field sprayer	Cucurbits Vegetables Crop Group 9; Grapes, Caneberries; Gooseberries; Saskatoon berries; Tomatoes, outdoor ornamentals		1	0	1	0
Airblast	Cucurbits Vegetables Crop Group 9; Grapes; Caneberries; Gooseberries; Saskatoon berries; Outdoor ornamentals	Early growth stage	1	0	1	0
		Late growth stage	1	0	1	0

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

ROTATIONAL CROP RESTRICTIONS

Crops on this label may be planted immediately after the last treatment. Do not plant cereal grains within 1 year after the last application. With the exception of cereal grains, crops not registered on this label may be planted 30 days after the last application.

INTEGRATED PEST MANAGEMENT

PROPERTY 300SC Fungicide is a disease control agent when used according to label directions for control of labeled fungi. PROPERTY 300SC Fungicide is recommended for use as part of an Integrated Pest Management (IPM) program, which may include the use of disease-resistant crop

varieties, cultural practices, crop rotation, biological disease control agents, pest scouting and disease forecasting systems aimed at preventing economic pest damage. Practices known to reduce disease development should be followed. Consult your provincial cooperative extension service or local agricultural authorities for additional IPM strategies established in your area. PROPERTY 300SC Fungicide may be used in disease forecasting programs that recommend application timing based upon environmental factors that favor disease development.

RESISTANCE MANAGEMENT

For resistance management, PROPERTY 300SC Fungicide contains a Group 50 fungicide. Any fungal population may contain individuals naturally resistant to PROPERTY 300SC Fungicide and other Group 50 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of PROPERTY 300SC Fungicide or other Group 50 fungicides with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.
- Fungicide use should be based on an integrated disease management (IPM) program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications.
- Monitor treated fungal populations for resistance development. Notify ISK Biosciences Corporation if reduced sensitivity of the pathogen to PROPERTY 300SC Fungicide is suspected.
- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information and to report suspected resistance, contact ISK Biosciences Corporation, at 1-877-706-4640

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.