



InCa™

Fertilizer with 5% Calcium

Guaranteed Minimum Analysis

Calcium (Ca).....5%

CAUTION

**AVOID FREEZING. WARM, SHAKE AND
RESUSPEND BEFORE USING**

KEEP OUT OF REACH OF CHILDREN

Belchim Crop Protection Canada, Inc.
104 Cooper Drive, Unit 3
Guelph, ON
N1C 0A4
519-826-7878

Manufactured by:
Plant Impact Plc
12 South Preston Office Village,
Cuerden Way, Bamber Bridge
Preston PR5 6BL United Kingdom

Net Contents: 10 L
Net Weight: 13 kg

General Information:

INCA can be applied to all crops including but not limited to potatoes, vegetable crops, grapes, melons, deciduous fruit trees, strawberries, cranberries, caneberries, cucurbits, lettuce and other leafy salads, tobacco and ornamental crops.

Note: Foliar fertilization is intended to supplement a standard ground fertility program. **INCA** should only be used on the basis of a soil and/or tissue analysis.

Do not combine with fertilizers that contain high levels of phosphate unless the specific combination is known to be compatible. Where no information is available on compatibility, it is recommended that a jar test be conducted.

Directions for Use

Apply **INCA** at 1.0 L/ha as a foliar spray or 1.5 L/ha as a foliar spray if water volumes exceed 600 litres per Ha

Method of Application

Conventional Sprayer: Apply the recommended amount of **INCA** in a minimum of 100L of water per hectare.

Aerial or Concentrate: Usually 50 L of water/ha is sufficient to apply the recommended amount of **INCA**. If lower volumes are used, a slight risk of foliage burning may occur on sensitive species.



Application Timing

Crop	Application Timing	
	Applications begin at:	Application Intervals
Potatoes	Tuber initiation	14 days
Apple, Stone and Tree Fruit	Early flower to harvest	7-14 days
Cabbage	3-5 leaf stage	14 days
Carrots, parsnips and sugar beets	3 leaf stage	14 days
Cucurbits, Tomatoes, Peppers, Melons	1 st flower to harvest	7-14 days
Strawberries and small fruit	Early bloom through harvest	14 days
Grape	Pre-bloom to harvest	14 day
Brassica Vegetables	3-5 leaf stage	14 days
Celery	3-5 leaf stage	14 days
Leafy Vegetables	3-5 leaf stage	14 days
Tobacco (flue cured and burley)	8-10 inch growth	14 days
Ornamentals	During periods of rapid growth	7-14 days

For full listings on application rates/timings please contact your Engage advisor.

Precautions

Product should be stored in the original container at ambient temperatures between 5 and 50°C. Store away from children, pets, livestock and foodstuffs. Avoid freezing and excessive temperature.

Wear gloves, eye protection and overalls when using the product. Avoid contact with skin and eyes. In case of contact with eyes, rinse immediately with plenty of water. Wash hands with soap and water before meals and after work.

Use this product in accordance with good agricultural practices. Do not make applications when temperatures are too hot, as crop damage may occur. Applications should be made at temperature levels and when other environmental conditions in your area are such that you experience indicates the application will be compatible and will accomplish the desired result.

Seller's Guarantee: Belchim Canada warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition. This product contains Plant Impact Technology

™ **INCA** is a trademark of Belchim Crop Protection Canada, Inc.