SLUGGO®
PROFESSIONAL



SLUGGO® PROFESSIONAL

Ferric phosphate (iron phosphate) is the active ingredient in Sluggo® Slug and Snail Bait, a selective molluscicide used to control slugs and snails in fruit, vegetable, ornamental, and greenhouse crops. Ferric phosphate is a naturally occurring iron compound that is formulated into attractive bait granules that slugs and snails consume while feeding. Once ingested, the active ingredient disrupts digestive and metabolic processes within the pest, causing it to stop feeding almost immediately. The pests then become less active, retreat to sheltered areas, and die over the following several days, often without leaving visible slime trails or dead bodies on the crop.

Key features and benefits of Sluggo® include effective slug and snail control with an excellent environmental and safety profile. Sluggo provides immediate crop protection because feeding ceases soon after bait consumption, reducing further plant damage even before the pests die. The bait is rain-resistant, easy to apply, requires no spraying or mixing, and remains effective under a wide range of environmental conditions. Sluggo phosphate is approved for use in organic production systems and is considered safe for use around crops, pets, wildlife, birds, and many beneficial organisms when used according to label directions. These attributes make Sluggo a valuable tool for protecting crop quality, reducing feeding damage, and managing slug and snail populations in both conventional and organic production systems.

Active Ingredient:

Iron 0.28% (present as Ferric Phosphate)

Key Crops:

Berries, vegetables, tree fruits, grapes, field crops, greenhouse fruit and vegetables, ornamentals

Key Targets:

Slugs, snails

IRAC Group:

N/A

Packaging Size:

20kg

Belchim Canada
GROWING TOGETHER

CROP	PEST	PHI	REI	RATE	ADDITIONAL INFORMATION
Root and Tuber Vegetables	Slugs, snails	0 days	4 hr.	12 - 50 kg/ha	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. At the seedling, transplant and later stages, apply the bait between the rows and around the perimeter of the field.
Bulb Vegetables	Slugs, snails	0 days	4 hr.	12 - 50 kg/ha	Use broadcast or granular spreader to apply evenly around or over the plants to be protected.
Leafy Vegetables	Slugs, snails	0 days	4 hr.	12 - 50 kg/ha	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. At the seedling, transplant and later stages, apply the bait between the rows and around the perimeter of the field.
Brassica Leafy Vegetables	Slugs, snails	0 days	4 hr.	12 - 50 kg/ha	
Legume Vegetables	Slugs, snails	0 days	4 hr.	12 - 50 kg/ha	
Fruiting Vegetables	Slugs, snails	0 days	4 hr.	12 - 50 kg/ha	
Cucurbit Vegetables	Slugs, snails	0 days	4 hr.	12 - 50 kg/ha	
Pome Fruit	Slugs, snails	0 days	4 hr.	12 - 50 kg/ha	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. For saplings, bait can be applied selectively in a band 15 cm wide around the base of the stem.
Stone Fruit	Slugs, snails	0 days	4 hr.	12 - 50 kg/ha	
Berries	Slugs, snails	0 days	4 hr.	12 - 50 kg/ha	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. The bait can be applied selectively in a 15cm wide band around the base of berry bushes.
Tree Nuts	Slugs, snails	0 days	4 hr.	12 - 50 kg/ha	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. For saplings, bait can be applied selectively in a band 15 cm wide around the base of the stem.
Cereal Grains	Slugs, snails	0 days	4 hr.	12 - 50 kg/ha	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. At the seedling and later stages, apply the bait between the rows and around the perimeter of the field.
Herbs and Spices	Slugs, snails	0 days	4 hr.	12 - 50 kg/ha	Use broadcast or granular spreader to apply evenly around or over the plants to be protected.
Oilseeds	Slugs, snails	0 days	4 hr.	12 - 50 kg/ha	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. At the seedling and later stages, apply the bait between the rows and around the perimeter of the field.
Stalk, Stem and Leaf Petioles	Slugs, snails	0 days	4 hr.	12 - 50 kg/ha	Use broadcast or granular spreader to apply evenly over the area.

Always read and follow the label.

Sluggo® Professional is a registered trademark of W. Neudorff GmbH KG

Belchim Canada
GROWING TOGETHER

519-826-7878
belchimcanada.com
104 Cooper Dr. Unit 3
Guelph, Ontario, Canada