

Beloukha®



BELOUKHA® HERBICIDE

Beloukha is a non-selective, bio-degradable, broad-spectrum, foliar-applied herbicide that acts exclusively by contact, attacking and destroying the cell membranes of the plant epidermis causing rapid tissue dehydration.

Beloukha is derived from a naturally occurring substance that is sourced from sunflowers and breaks down into carbon dioxide and water. Beloukha provides burndown of both annual and perennial broadleaf and grass weeds, plus most mosses and other cryptogams (liverworts, lichens, algae).

| |
|---|
| Active Ingredient: |
| Pelargonic Acid 500 g / L |
| Key Registered Crops: |
| Fruit & Vegetable crops, Cereal crops, Potatoes |
| HRAC Group: |
| Group 0 |
| Packaging Size: |
| 2 x 9.46L |

APPLICATION INFORMATION FOR BELOUKHA®

| Crop | Purpose | Application | REI | PHI | Rate |
|---|--|--|----------|-------|---|
| Berries & small fruit Stone fruit Pome fruit Fruiting vegetables Cucurbits Root & Tuber veg. Legumes Cereal grains | Weed control | Use minimum effective rate for weed control. For harder to control weeds, higher rates or repeat applications may be required. Use shielded / directed spray to avoid spraying desired vegetation. Use on woody / hardened vines or orchard trees that are more than 2 years old | When dry | 1 day | 16 - 27 L / ha in 200 – 300 L / ha water |
| Potato | Desiccation | For use with or without mechanical top beating. Use lower application rates when using mechanical top beater and use higher rates without. | When dry | 1 day | 16 - 22 L / ha in 300 L / ha water |
| Wheat, Barley, Oats | Desiccation; Pre-harvest weed suppression | Apply at onset of senescence and harvest when desired moisture level is reached. | When dry | 1 day | 11 - 22 L / ha in 200 – 300 L / ha water |
| Grapes Stone fruit Pome fruit | Suckers | Use shielded / directed spray to avoid spraying desired vegetation. Use on woody / hardened vines or orchard trees older than 2 years old. Higher rates or repeat applications may be required for larger sized suckers. | When dry | 1 day | 4 - 7 L / ha in 100 – 200 L / ha water |
| Post-harvest | Weeds | Use shielded / directed spray to avoid spraying desired vegetation. For hard to control weeds, higher rates or repeat applications may be required. | When dry | NA | 16 - 27 L / ha in 200 – 300 L / ha water |

Always read and follow the label. Do not make more than 2 applications per season.

- Contact, non-selective, broad spectrum, foliar applied herbicide
- Low odour through improved formulation
- Biodegradable
- Provides burndown of both annual and perennial broadleaf weeds, grasses and more
- Fast acting with visible effects on most weeds within hours of application
- Versatile - for use on a wide variety of crops