

Container

Kenja[®] 400SC Fungicide

COMMERCIAL

SUSPENSION

READ THE LABEL AND BOOKLET BEFORE USING

ACTIVE INGREDIENT: Isofetamid 400 g/L

Contains 1,2-benzisothiazolin-3-one at 0.017% as a preservative

OR

Contains 2-Bromo-2-nitropropane-1,3-diol at 0.0079%

and

5-Chloro-2-methyl-4-isothiazolin-3-one at 0.00076%

and

2-Methyl-4-isothiazolin-3-one at 0.00025% as preservatives

REGISTRATION NO: 31758 PEST CONTROL PRODUCTS ACT

Net Contents: 500 ml to 200 L

Registrant
ISK Biosciences Corporation
7470 Auburn Road, Suite A
Concord, Ohio
44077
USA
1-877-706-4640

Distributed By:
Belchim Crop Protection
104 Cooper Drive, Unit 3
Guelph, ON
N1C 0A4
Canada
866-613-3336

Kenja is a registered trademark of Ishihara Sangyo Kaisha, Ltd.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing/loading, application and during clean-up and repair. Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Do not use in residential areas. Residential areas are defined as any use site where bystanders including children could be exposed during or after application. This includes homes, schools, parks, playgrounds, playing fields, public buildings or any other area outside of golf courses where the general public could be exposed.

Tank Mixes: When applied as a tank-mix combination, read and observe all label precautionary measures, including personal protection equipment, directions for use, application rates, and restrictions for each product used in the tank-mix. Follow the more stringent statements on the product labels for mixing, loading, applying, and for post-application entry into treated areas.

DO NOT allow releases, effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters..

Do not apply by air.

FIRST AID

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so

by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

In Case of Emergency: Call 1-888-484-7546.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

Toxic to birds, small wild mammals and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.”

STORAGE

Store this product away from food and feed.

If this product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada’s web site at www.croplife.ca.

DISPOSAL

Recyclable containers

Disposal of container: DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray

- mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Disposal of unused, unwanted product: For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Pamphlet

Kenja[®] 400SC Fungicide

COMMERCIAL

SUSPENSION

READ THE LABEL AND BOOKLET BEFORE USING

ACTIVE INGREDIENT: Isofetamid 400 g/L

Contains 1,2-benzisothiazolin-3-one at 0.017% as a preservative

OR

Contains 2-Bromo-2-nitropropane-1,3-diol at 0.0079%

and

5-Chloro-2-methyl-4-isothiazolin-3-one at 0.00076%

and

2-Methyl-4-isothiazolin-3-one at 0.00025% as preservatives

REGISTRATION NO: 31758 PEST CONTROL PRODUCTS ACT

Net Contents: 500 ml to 200 L

Registrant
ISK Biosciences Corporation
7470 Auburn Road, Suite A
Concord, Ohio
44077
USA
1-877-706-4640

Distributed By:
Belchim Crop Protection
104 Cooper Drive, Unit 3
Guelph, ON
N1C 0A4
Canada
866-613-3336

Kenja is a registered trademark of Ishihara Sangyo Kaisha, Ltd.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing/loading, application and during clean-up and repair. Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Do not use in residential areas. Residential areas are defined as any use site where bystanders including children could be exposed during or after application. This includes homes, schools, parks, playgrounds, playing fields, public buildings or any other area outside of golf courses where the general public could be exposed.

Tank Mixes: When applied as a tank-mix combination, read and observe all label precautionary measures, including personal protection equipment, directions for use, application rates, and restrictions for each product used in the tank-mix. Follow the more stringent statements on the product labels for mixing, loading, applying, and for post-application entry into treated areas.

DO NOT allow releases, effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters..

Do not apply by air.

FIRST AID

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so

by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

In Case of Emergency: Call 1-888-484-7546.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

Toxic to birds, small wild mammals and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.”

STORAGE

Store this product away from food and feed.

If this product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada’s web site at www.croplife.ca.

DISPOSAL

Recyclable containers

Disposal of container: DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking

the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Disposal of unused, unwanted product: For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

DIRECTIONS FOR USE

KENJA 400SC Fungicide is a broad-spectrum fungicide with preventative, local systemic (or translaminar) and curative properties for foliar and soil-borne diseases. KENJA 400SC Fungicide must be applied in scheduled protective programs and used in rotation with products with a different mode of action.

Under conditions favorable for disease development, the highest rate specified and shortest application interval should be used. Thorough, uniform coverage is essential for disease control.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

DO NOT apply by air.

MIXING

NOTE: Slowly invert container several times to assure uniform mixture of formulation before adding this product to the spray tank.

The required amount of KENJA 400SC Fungicide should be added slowly into the spray tank during filling. With concentrate sprays, pre-mix the required amount of KENJA 400SC Fungicide in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations. **DO NOT** allow spray mixture to stand overnight or for prolonged periods. Prepare only the amount of spray required for immediate use. Spraying equipment should be thoroughly cleaned immediately after the application.

TANK MIXING AND COMPATIBILITY

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

KENJA 400SC Fungicide is generally compatible with other fungicides, insecticides, fertilizers and micronutrient products provided sufficient free water is available for dispersion of all the tank mix products. However, the physical compatibility of this product with tank mix partners must be evaluated before use. Conduct a jar test with intended tank-mix product(s) prior to preparation of large volumes. Use the following procedure: 1) Pour the recommended proportions of the products into a suitable container of water, 2) Mix thoroughly and 3) Allow to stand 5 minutes. If the combination remains mixed or can be re-mixed readily, it is considered physically compatible. Any physical incompatibility in the jar test indicates that KENJA 400SC Fungicide must not be used in the tank-mix. Since not all crop varieties have been tested with all possible tank mix combinations, the combination should also be tested for crop safety on a small portion of the crop to ensure that phytotoxicity will not occur.

In some cases, tank mixing pest control products can result in reduced pest efficacy or increased host crop injury. The user should contact ISK Biosciences Corporation at 1-877-706-4640 for information before applying any tank mix that is not specifically recommended on this label.

APPLICATION

Apply KENJA 400SC Fungicide in sufficient water to obtain adequate coverage of the foliage. Water volume to be used will vary with crop and amount of plant growth. Spray volume will usually range from 200 to 1000 litres per hectare for dilute sprays, typically used in orchards or for large crops. For low growing crops a spray volume of 50 to 100 litres per hectare is recommended.

ROTATIONAL CROP RESTRICTIONS

Crops on this label may be planted immediately after the last treatment. Do not plant other crops not registered for this product within 30 days after the last application.

INTEGRATED PEST MANAGEMENT

KENJA 400SC Fungicide is an excellent disease control agent when used according to label directions for control of labeled fungi. KENJA 400SC Fungicide is recommended for use as part of an Integrated Pest Management (IPM) program, which may include the

use of disease-resistant crop varieties, cultural practices, crop rotation, biological disease control agents, pest scouting and disease forecasting systems aimed at preventing economic pest damage. Practices known to reduce disease development should be followed. Consult your provincial cooperative extension service or local agricultural authorities for additional IPM strategies established in your area. KENJA 400SC Fungicide may be used in disease forecasting programs that recommend application timing based upon environmental factors that favor disease development.

RESISTANCE MANAGEMENT

For resistance management, please note that KENJA 400SC Fungicide contains a Group 7 fungicide. Any fungal population may contain individuals naturally resistant to KENJA 400SC Fungicide and other Group 7 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of KENJA 400SC Fungicide or other Group 7 fungicides with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.
- Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant disease resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications.
- Monitor treated fungal populations for resistance development. Notify ISK Biosciences Corporation if reduced sensitivity of the pathogen to KENJA 400SC Fungicide is suspected.
- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information and to report suspected resistance, contact ISK Biosciences Corporation at 1-877-706-4640.

DIRECTIONS FOR USE

Crop	Diseases	Use Rate	Instructions
Grapes	Control of Botrytis bunch rot (<i>Botrytis cinerea</i>)	1.46 to 1.61 L/ha	<p>Application Instructions: For use on all types of grapes (wine, table, raisin, and juice).</p> <p>For bunch rot make applications at critical timings for Botrytis control on a 14 day interval. Applications are typically made at early bloom, bunch closure, veraison and pre-harvest. Apply with sufficient water to allow for penetration into the foliage to obtain complete coverage.</p> <p>Resistance Management: Do not make more than 2 sequential applications of KENJA 400SC Fungicide or other Group 7 containing fungicides before rotating to a fungicide with a different mode of action.</p> <p>Restrictions: Do not apply more than 3 applications/year The Pre-Harvest Interval (PHI) for this crop is 14 days.</p>
Lettuce Head and Leaf	Control of Sclerotinia drop (<i>Sclerotinia minor</i> , <i>Sclerotinia sclerotiorum</i>)	0.90 L/ha	<p>Application Instructions: On direct seeded lettuce make the first application after emergence, thinning or prior to onset of disease development</p> <p>On transplanted lettuce make the first application immediately after transplanting or prior to the onset of disease development.</p> <p>Make a second application 14 days later if conditions continue to favor disease development.</p> <p>Resistance Management: Do not make more than 2 sequential applications of KENJA 400SC Fungicide</p>

			<p>or other Group 7 containing fungicides before rotating to a fungicide with a different mode of action.</p> <p>Restrictions: Do not apply more than 2 applications/year The Pre-Harvest Interval (PHI) for this crop is 14 days.</p>
<p>Low Growing Berry, Crop Subgroup 13-07G</p>	<p>Control of Grey mold (<i>Botrytis cinerea</i>)</p>	<p>0.987 - 1.24 L/ha</p>	<p>Application Instructions: Initiate applications prior to disease development and continue on a 7- to 14-day interval. When disease pressure is high use the high rate and shortest interval.</p> <p>Resistance Management: Do not make more than 2 sequential applications of KENJA 400SC Fungicide or other Group 7 containing fungicides before rotating to a fungicide with a different mode of action.</p> <p>Restrictions: Do not apply more than 5 applications /year The Pre-Harvest Interval (PHI) for this crop is 0 days.</p>
<p>Includes all members of the Crop Subgroup 13-07G, Bearberry; bilberry; blueberry, lowbush; cloudberry; cranberry; lingonberry; muntries; partridgeberry; strawberry; and cultivars, varieties, and/or hybrids of these.</p>			
<p>Berry and Small Fruit Crop Group 13-07</p> <p>except Subgroup 13-07C, 13-07F, 13-07G. For instructions on</p>	<p>Control of Grey mold (<i>Botrytis cinerea</i>)</p>	<p>0.987 - 1.24 L/ha</p> <p>(395 – 496 g a.i./ha)</p>	<p>Application Instructions: Initiate applications prior to disease development and continue on a 7- to 14-day interval. When disease pressure is high use the highest rate and shortest interval.</p> <p>Resistance Management: Do not make more than 2 sequential applications of KENJA 400SC</p>

<p>grape and 13-07G, refer to tables above. DO NOT USE ON CROP SUBGROUP 13-07C.</p>			<p>Fungicide or other Group 7 containing fungicides before rotating to a fungicide with a different mode of action.</p> <p>Restrictions: Do not apply more than 3 applications /year. The Pre-Harvest Interval (PHI) for crop group is 7 days.</p>
<p>Includes all members of the Berry and Small Fruit group except Subgroups 13-07C, 13-07F and 13-07G: Aronia berry; blackberry; blueberry, highbush; Chilean guava; cranberry, highbush; currant (buffalo, black, red, native); European barberry; gooseberry; honeysuckle, edible; huckleberry; jostaberry; kiwifruit, fuzzy;raspberry, black and red; sea buckthorn; wild raspberry; and cultivars, varieties, and/or hybrids of these.</p>			
<p>Stone Fruit, Crop Group 12-09</p>	<p>Suppression of Brown Rot and Blossom Blight (<i>Monilinia fructicola</i>)</p>	<p>0.913 L/ha (365 g a.i./ha)</p>	<p>Application Instructions: Initiate applications prior to disease development and continue on a 7 to 14-day interval. Apply with a silicone surfactant, such as Xiameter OFX 0309 Fluid, at a rate of 0.1%.</p> <p>Resistance Management: Do not make more than 2 sequential applications of KENJA 400SC Fungicide or other Group 7 containing fungicides before rotating to a fungicide with a different mode of action.</p> <p>Restrictions: Do not apply more than 3 applications /year The Pre-Harvest Interval (PHI) for this crop group is 1 day.</p> <p>Do not apply in a spray volume greater than 1000 L/ha. A minimum concentration of 0.913 ml/L spray solution is required for adequate control.</p>
<p>Includes all members of the Stone Fruit Crop Group Family including apricot; apricot, Japanese; cherry, black; cherry, Nanking; cherry, sweet; cherry, tart; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these.</p>			

<p>Apple</p>	<p>Control of Apple Scab (<i>Venturia inaequalis</i>)</p>	<p>0.913 L/ha (365 g a.i./ha)</p>	<p>Application Instructions: Initiate applications prior to disease development and continue on a 10 to 14-day interval.</p> <p>Resistance Management: Do not make more than 2 sequential applications of KENJA 400SC Fungicide or other Group 7 containing fungicides before rotating to a fungicide with a different mode of action.</p> <p>Restrictions: Do not apply more than 6 applications /year The Pre-Harvest Interval (PHI) for this crop group is 20 days.</p> <p>Do not apply in a spray volume greater than 1000 L/ha.</p>
<p>Edible-podded Legume Vegetables, (Crop Subgroup 6A) except pigeon pea and pea (Pisum spp, which includes dwarf pea, edible-podded pea, snow pea, sugar snap pea)</p> <p>Succulent shelled pea and bean (Crop Subgroup 6B)</p>	<p>Suppression of White mold, (<i>Sclerotinia sclerotiorum</i>)</p>	<p>1.25 L/ha (500 g a.i./ha)</p>	<p>Application Instructions: Initiate applications prior to disease development and continue on a 7 to 14-day interval. Beans and other legumes are most susceptible to white mold infection during the bloom period.</p> <p>Resistance Management: Do not make more than 2 sequential applications of KENJA 400SC Fungicide or other Group 7 containing fungicides before rotating to a fungicide with a different mode of action. When applied to soybean (immature seed), for resistance management purposes, Kenja 400SC should be tank-mixed with Allegro 500F Agricultural Fungicide. Consult the Allegro 500F label for additional instructions and precautions.</p> <p>Restrictions: Do not apply more than 2 applications /year</p>

Dried shelled pea and bean except soybean (Crop Subgroup 6C)			<p>The Pre-Harvest Interval (PHI) for edible-podded snap beans is 7 days.</p> <p>The Pre-Harvest Interval (PHI) for succulent shelled peas and beans is 14 days.</p> <p>The PHI for dried shelled peas and beans is 30 days.</p>
<p>Edible-podded Legume Vegetables Subgroup 6A: bean (<i>Phaseolus spp.</i>) (includes runner bean, snap bean, wax bean); bean (<i>Vigna spp.</i>) (includes asparagus bean, Chinese longbean, moth bean, yardlong bean); jackbean; soybean (immature seed); sword bean</p> <p>Succulent Shelled Pea and Bean Subgroup 6B: bean (<i>Phaseolus spp.</i>) (includes lima bean, green; bean (<i>Vigna spp.</i>) (includes blackeyed pea, cowpea, southern pea); broad bean (fava bean) (<i>Vicia faba</i>); pea (<i>Pisum spp.</i>) (includes English pea, garden pea, green pea); pigeon pea (<i>Cajanus cajan</i>)</p> <p>Dry Shelled Pea and Bean except Soybean Subgroup 6C: dried cultivars of bean (<i>Lupinus spp.</i>) (includes grain lupin, sweet lupin, white lupin and white sweet lupin) ; bean (<i>Phaseolus spp.</i>) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); bean (<i>Vigna spp.</i>) (includes adzuki bean, blackeyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); broad bean (fava bean - dry) (<i>Vicia faba</i>); chickpea (garbanzo bean); guar; lablab bean (hyacinth bean); lentil; pea (<i>Pisum spp.</i>) (includes field pea); pigeon pea</p>			

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for this product for the use(s) described below were developed by persons other than ISK Biosciences Corporation under the User Requested Minor Use Label Expansion program. ISK Biosciences Corporation has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

DIRECTIONS FOR USE

Crop	Diseases Controlled	Use Rate	Instructions
Ginseng	<p><i>Cylindrocarpon</i> root rot</p> <p>(Rusty Root Disease)</p> <p>(<i>Ilyonectria mors-panacis</i> syn. <i>Cylindrocarpon destructans</i>)</p>	<p>1.46 L product/ha</p> <p>(584 g a.i./ha)</p>	<p>Application Instructions: Initiate applications within 2 weeks of plant emergence and/or prior to disease development and continue at 7-day intervals if conditions continue to be favorable for disease development.</p> <p>Use 1000 to 1870 L/ha of spray volume to ensure runoff into the soil and root zone.</p> <p>Resistance Management: Do not make more than 2 sequential applications of KENJA 400SC FUNGICIDE or other Group 7 containing fungicides before rotating to a fungicide with a different mode of action.</p> <p>Restrictions: Do not apply more than 3 applications per year. The Pre-Harvest Interval (PHI) for this crop is 14 days.</p>
Greenhouse ornamental plants	Botrytis grey mould (<i>Botrytis cinerea</i>)	78 to 133 mL Product Per 100 L of Water	<p>Application Instructions: Apply as a foliar broadcast spray. Applications should begin when conditions are favorable for disease development or when plants first exhibit disease symptoms, from germination to mature crop. Within the stated rate range, use the lower rate and shortest interval and the higher rate at the longest interval. When disease is severe use the highest labelled rate at the shortest labelled interval.</p> <p>Repeat applications on a 7-14 day interval.</p> <p>The spray volume should generally not exceed 1000 L per hectare. If a larger volume is needed to wet the foliage, do not</p>

			<p>exceed the maximum rate of 1333 ml product/ha.</p> <p>Resistance Management: Do not make more than 2 sequential applications of KENJA 400SC FUNGICIDE or other Group 7 containing fungicides before rotating to a fungicide with a different mode of action.</p> <p>Restrictions: Do not apply more than 3 applications at the highest rate per crop cycle or 6 applications at the lowest rate per crop cycle.</p>
--	--	--	--

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Airblast application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. DO NOT apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

Spray Buffer Zones:

Use of the following spray methods or equipment DO NOT require a spray buffer zone: hand-held or backpack sprayer and spot treatment.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Spray Buffer Zones (metres) Required for the Protection of:			
			Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:	
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m
Field sprayer	Lettuce, Oilseed Crop Group 20A Edible-podded Legume Vegetables (Crop Subgroup 6A), Succulent shelled pea and bean (Crop Subgroup 6B, Dried shelled pea and bean except soybean (Crop Subgroup 6C) Berry and Small Fruit (Crop 13-07 except Subgroups 13-07C, 13-07F and 13-07G), Ginseng		1	0	0	0
	Low-growing berries (Crop Group 13-07G)		1	1	1	0
	Turf		2	1	1	0
Airblast	Grapes, Pome Fruit (Crop Group 11-09)	Early growth stage	15	1	1	0
		Late growth stage	5	1	1	0
	Low-growing berries (Crop Group 13-07G)	Early growth stage	15	1	1	0
		Late growth stage	10	1	1	0
	Berry and Small Fruit (Crop 13-07 except Subgroups 13-07C, 13-07F and 13-07G)	Early growth stage	10	0	0	0
		Late growth stage	5	0	0	0
	Stone Fruit (Crop Group 12-09)	Early growth stage	10	0	0	0
		Late growth stage	4	0	0	0

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.